



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07F 9/50, B01J 31/24, C07C 67/36	A1	(11) International Publication Number: WO 99/47528 (43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/GB99/00797 (22) International Filing Date: 16 March 1999 (16.03.99) (30) Priority Data: 9805348.1 16 March 1998 (16.03.98) GB (71) Applicant (for all designated States except US): IMPERIAL CHEMICAL INDUSTRIES PLC [GB/GB]; Imperial Chem- ical House, Millbank, London SW1P 3JF (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): NEWMAN, Paul, David [GB/GB]; 20 Saint Margarets Court, Linnet Close, Cyncoed, Cardiff CF2 7HG (GB). CAMPBELL, Richard, Anthony [GB/GB]; 4 Tulloch Place, Saint Andrews, Fife KY16 8XJ (GB). TOOZE, Robert, Paul [GB/GB]; 11 Station Road, Norton, Stockton, Cleveland TS20 1EA (GB). EASTHAM, Graham, Ronald [GB/GB]; 7 Heslop Drive, Darlington, County Durham DL1 5TQ (GB). THORPE, Jamie, Michael [GB/GB]; 19 Woodside, Redcar, Cleveland TS10 4NG (GB). EDWARDS, Peter, Gerald [GB/GB]; 81 Plymouth Road, Penarth, Cardiff CF64 3DE (GB).		(74) Agents: GIBSON, Sara, Hillary, Margaret et al.; ICI Group Intellectual Property, Wilton, Middlesbrough, P.O. Box 90, Cleveland TS90 8JE (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROCESS FOR THE PREPARATION OF BISPHOSPHINES		
(57) Abstract <p>A method of manufacturing a compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$, in which each R is independently a pendant, optionally substituted, organic group through which the group is linked to tertiary carbon atom C; L^1, L^2 are independently a linking group selected from an optionally substituted lower alkylene chain connecting the respective phosphorus atom to the group X and X is a bridging group comprising an optionally substituted aryl moiety to which the phosphorus atoms are linked on available adjacent carbon atoms, comprises: i) reacting together a compound of formula $H-L^1-X-L^2-H$ with an organometallic compound to form an intermediate compound of formula $M-L^1-X-L^2-M$, where M is an alkali metal atom; ii) reacting said intermediate compound with a compound of formula $(R_3-C)_2P-A$, where A is a halogen atom, to form said compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$. M is preferably lithium, potassium or sodium and the intermediate compound may be isolated or not. The reaction (i) may advantageously be carried out in the presence of a complexing agent such as tetramethylethylenediamine.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PROCESS FOR THE PREPARATION OF BISPHOSPHINES

The present invention relates to a phosphine compound which is useful as a component of a catalyst system which may be used in the carbonylation of olefins, and in particular to a method of manufacturing such phosphine compounds.

- 5 WO 96/19434 discloses a process for the carbonylation of ethylene and a catalyst system for use therein. The catalyst system described in that document comprises a bidentate phosphine of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$, in which each R is independently a pendant, optionally substituted, organic group through which the group is linked to tertiary carbon atom C; L^1 , L^2 are independently a linking group selected from an
- 10 optionally substituted lower alkylene chain connecting the respective phosphorus atom to the group X and X is a bridging group comprising an optionally substituted aryl moiety to which the phosphorus atoms are linked on available adjacent carbon atoms. One example of such a bidentate phosphine is bis (di-*t*-butyl phosphino)-*o*-xylene (also known as 1,2-bis(di-*t*-butylphosphinomethyl)benzene).
- 15 Such catalysts may be made by mixing the phosphine ligand with a suitable source of palladium such as palladium acetate. WO 96/19434 describes the preparation of one form of the ligand via the phosphonium salt produced from the reaction of the appropriate secondary phosphine with the corresponding aromatic dihalide. In the preferred form of the phosphine ligand in WO 96/19434, R is a lower alkyl group, in particular methyl. A
- 20 problem with manufacturing this ligand by the route described is that the secondary phosphine which is used (e.g. di-*t*-butyl phosphine) is toxic, highly reactive, smelly and flammable. We have also found that the reaction is low yielding and converts some of the di-*t*-butyl phosphine to a non-reclaimable waste product which must be disposed of.
- Al-Salem et al in Journal of the Chemical Society (Dalton) 1979 page 1980 describes
- 25 making 1,5 bis(di-*t*-butylphosphino)pentane by reacting lithium metal with 1,5-dibromopentane and then phosphorylating the resulting lithiated intermediate with *t*-butylchlorophosphine. This method, for forming a phosphine of an alkyl compound, starts from the halogenated alkyl compound. Alkyl halides typically require considerable care during storage and use and may be very unpleasant to use. Therefore using this
- 30 method of making the phosphine using the non-halogenated alkyl compound as a starting material would require an extra step of converting first to the alkyl halide.

We have now found that phosphine ligands of the type described in WO 96/19434 may be prepared by a high yielding route using more benign materials which produce less waste phosphorus product than the route described in WO 96/19434.

According to the invention, a method of manufacturing a compound of general formula
 5 $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$, in which each R is independently a pendant, optionally substituted, organic group through which the group is linked to tertiary carbon atom C; L^1 , L^2 are independently a linking group selected from an optionally substituted lower alkylene chain connecting the respective phosphorus atom to the group X and X is a bridging group comprising an optionally substituted aryl moiety to which the phosphorus atoms are
 10 linked on available adjacent carbon atoms comprises the steps of
 i) reacting together a compound of formula $H-L^1-X-L^2-H$ with an organometallic compound to form an intermediate compound of formula $M-L^1-X-L^2-M$, where M is an alkali metal atom,
 ii) reacting said intermediate compound with a compound of formula $(R_3-C)_2P-A$, where A
 15 is a halogen atom, to form said compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$.

The compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$ may be useful as a component of a catalyst compound. In particular WO 96/19434 describes the use of such compounds as bidentate ligands which, when used together with a Group VIII metal such as palladium, are effective catalysts for the carbonylation of olefins.

20 The pendant, optionally substituted organic groups, R may be independently selected from a wide range of components. Preferably, the pendant groups are optionally substituted lower alkyl, e.g. C_{1-8} , and may be branched or linear.

Particularly preferred is when the organic groups, R, when associated with their respective
 25 carbon atom, form composite groups which are at least as sterically hindering as *tert*-butyl. Steric hindrance in this context is as discussed at page 14 et seq of "Homogeneous Transition Metal Catalysis - A Gentle Art", by C Masters, published by Chapman and Hall, 1981. In one preferred embodiment, the organic groups R are all methyl groups, i.e. a preferred form of R_3-C group is a tertiary butyl group.

30 The linking groups, L^1 and L^2 , are independently selected from an optionally substituted,

particularly lower alkyl, e.g. C₁ to C₄, substituted, lower alkylene, e.g. C₁ to C₄ chain. Especially preferred is when both L¹ and L² are methylene.

The bridging group X is an aryl moiety, e.g. a phenyl group, which may be optionally substituted, provided that the two phosphorus atoms are linked to adjacent carbon atoms, e.g. at the 1 and 2 positions on the phenyl group. Optional substitution of the aryl moiety may be by other organic groups, e.g. alkyl, particularly C₁₋₈, aryl, alkoxy, carbalkoxy, halo, nitro, trihalomethyl and cyano. Furthermore, the aryl moiety may be a fused polycyclic group, e.g. naphthalene, biphenylene or indene.

10

Examples of compounds which may advantageously be made by the method according to the invention are bis (di-*tert*-butyl phosphino) - o - xylene (also known as 1,2 bis (di-*tert*-butylphosphinomethyl) benzene), bis (di-*t*-neopentyl phosphino) - o - xylene and bis 1, 2 (di-*tert*-butyl phosphino) naphthalene.

15 The reaction of a compound of formula H-L¹-X-L²-H with a basic organometallic compound of formula R^mM to form an intermediate compound of formula M-L¹-X-L²-M, where M is an alkali metal atom, may be carried out by various means which are known in the art of organometallic chemistry. For example such metallation methods are described by Wilkinson *et al* in "Comprehensive Organometallic Chemistry" at page 54; and by Lambert
20 *et al* in "Preparative Polar Organometallic Chemistry".

The organometallic compound may comprise a compound of formula R^m-M, where R^m is an organic group which tends to withdraw electrons from the metal atom M. Suitable organic groups include aromatic or aliphatic groups, especially alkyl groups, which may be substituted. Lower alkyl groups have been found to be particularly suitable, for example
25 preferred R^m compounds include n-butyl, *t*-butyl, *sec*-butyl, methyl or pentyl. M may be any suitable alkali metal which forms a polar organometallic group with R^m. Suitable metals include those of Group IA, e.g. sodium, potassium or lithium. When M comprises K or Na the R^m-M metallating agent is preferably generated in situ by an exchange mechanism, e.g. by the reaction between R^m-Li and potassium or sodium
30 *t*-butoxide as described by Lochman *et al* in Tetrahedron Letters No. 2 pages 257 - 262 (1966). Preferred metallating compounds are butyl lithium, butyl sodium and butyl potassium, the latter compounds preferably being formed in situ by the reaction of butyl lithium with potassium or sodium *t*-butoxide.

Alternative organometallic compounds are also known in the art and may comprise $\text{Me}_3\text{SiCH}_2\text{K}$, alkali amides MNH_2 preferably used in liquid ammonia, lithium dialkylamides e.g. lithium diisopropylamide (LDA), lithium, sodium or potassium metals, metal hydrides e.g. KH in the presence of co-ordinating compounds.

- 5 The reaction between R^mM and $\text{H-L}^1\text{-X-L}^2\text{-H}$ may be carried out in the presence of a solvent. Any solvent used must not contain any component which reacts with the intermediate compound, and suitable such solvents will be well known to the skilled chemist. Favoured solvents include dry alkyl ethers e.g. diethyl ether, methyl t-butyl ether, di(n-propyl) ether; tetrahydrofuran (THF), and hydrocarbons such as hexane and heptane.
- 10 The reaction between R^mM and $\text{H-L}^1\text{-X-L}^2\text{-H}$ may be beneficially carried out in the presence of a basic compound which is capable of forming a complex with the metal. A preferred complexing agent is tetramethylethylenediamine (TMEDA). The presence of TMEDA is greatly preferred when the metallating agent is alkyl lithium or alkyl sodium. When alkyl potassium (or alkyl lithium/ potassium t-butoxide mixture) is used, we have
- 15 found that the reaction proceeds satisfactorily in the absence of TMEDA.

The mole ratios of metallating agent : $\text{H-L}^1\text{-X-L}^2\text{-H}$ are preferably in the range 1:1 to 10:1, more preferably at least 1.5:1. The preferred ratio depends upon the nature of the metallating agent used, for example when alkyl lithium is used a ratio of $\text{R-Li} : \text{H-L}^1\text{-X-L}^2\text{-H}$ of 3:1 may be preferred. Higher ratios may be preferred to encourage the formation of

20 di-substituted product rather than mono-substituted compounds.

A preferred ratio of $\text{R}^m\text{M} : \text{complexing agent}$ is in the range 1:1 - 4:1, especially preferred is a ratio of about 1:1 to about 2:1.

- It is preferred to conduct the metallation reaction between R^mM and $\text{H-L}^1\text{-X-L}^2\text{-H}$ at a temperature in the range -20 to 150 °C, more preferably at room temperature or above.
- 25 The optimum temperature and reaction time depends upon the identity of the reactants, in particular upon the alkali metal which is used. For example we have found, using o-xylene as the compound to be metallated that when alkyl lithium is used, the reaction proceeds well at room temperature (i.e. about 20 - 22 °C) whereas reactions using alkyl

sodium and potassium are preferably conducted at about 60, (e.g. 50 - 70 °C) and about 80 °C (e.g. 70 - 90 °C) respectively.

The intermediate compound, $M-L^1-X-L^2-M$ may be isolated from the reaction mixture prior to conducting the reaction between $M-L^1-X-L^2-M$ and the compound of formula

- 5 $(R_3-C)_2P-A$. However isolation of the intermediate may not be necessary and the reaction has been found to proceed very satisfactorily when the intermediate is not isolated. When the intermediate product $M-L^1-X-L^2-M$ can be separated relatively easily, for example if it is in a different physical form from the starting materials, then it may be advantageous to isolate the intermediate from the reaction mixture to encourage the formation of more
- 10 intermediate. We have found that the reaction mixture may contain partially reacted compounds of formula $M-L^1-X-L^2-H$ which may be reacted to completion by the addition of further quantities of R^mM and complexing agent if used.

- The phosphorylation reaction between the intermediate product $M-L^1-X-L^2-M$ and the halophosphine $(R_3-C)_2P-A$ is also carried out in the presence of a solvent and similar
- 15 solvents to those used for the metallation are suitable, e.g. ethers, C5+ alkanes and petroleum ethers. We have found that, when the metal used is potassium, the reaction is favoured when diethyl ether is used as a solvent for the phosphorylation reaction, although an alkane e.g. heptane, may be preferred for the metallation reaction. Also, even when sodium or lithium is used as the metal species, it may be beneficial to use a
- 20 different solvent for the phosphorylation from that used for the metallation, e.g. to enable higher temperatures to be used in one reaction yet allow an easy separation of the solvent from the product of the second reaction.

- The phosphorylation reaction may be conducted at elevated temperatures, e.g. at temperatures of 60°C or greater, but it is preferred to conduct the reaction at room
- 25 temperature or below, e.g. at -20 to 25 °C.

The compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$ may be isolated from the reaction mixture by distilling off the excess solvent, preferably under vacuum and then extracting the product compound into a solvent, e.g. methanol, from which it may be precipitated.

The invention will be further described, by way of example only, below. In all reactions, the reactants and apparatus used were prepared to allow the reaction to proceed in anhydrous and anaerobic reaction conditions.

Example 1

- 5 *N,N,N',N'*-Tetramethylethylenediamine (0.79cm³, 5.25 mmol) was added to a stirred solution of *n*-butyllithium (2.10cm³, 2.5M in hexane, 5.25 mmol) in heptane (15cm³) under an atmosphere of nitrogen. After five minutes, *ortho*-xylene (0.26cm³, 2.10mmol) was added slowly. A orange-brown precipitate formed after leaving the solution stirring at room temperature for 96 hours.
- 10 Di-*tert*-butylchlorophosphine (1.40cm³, 7.37 mmol) was added, then at room temperature the solution was left to stir until the colour disappeared.

A sample of the reaction mixture was analysed by ³¹P{¹H} NMR and the conversion to bis (di-*tert*-butyl phosphino)-*o*-xylene and to the mono-substituted analogue, was determined. The results are shown in Table 1.

15 Examples 2 - 8

The experimental procedure of Example 1 was repeated but varying the reactant ratios and reaction conditions as shown in Table 1.

Example 9

- 20 The experimental procedure described above was repeated using a ratio of butyllithium:TMEDA: *o*-xylene of 4 : 4 : 1 , a temperature of 25°C and a reaction time of 48 hrs, but the product of the reaction between the butyl lithium and *o*-xylene / TMEDA was filtered to separate the brown solid from the red liquid. Each was then reacted separately with di-*tert*-butylchlorophosphine and the conversion to bis (di-*tert*-butyl phosphino) - *o* - xylene and to the mono-substituted analogue, was determined. The results are shown in Table
- 25 2.

Examples 10 - 13

n-Butyllithium (11 mmol) was added dropwise to a well-stirred solution of *o*-xylene (0.6 ml, 5 mmol) and potassium *tert*-butoxide (1.2g 11mmol) in heptane (25 ml). The reactions were stirred (and sometimes heated) for a predetermined length of time (see Table 3).

The system was cooled by placing it in an ice bath and di-*tert*-butylchlorophosphine (1.87 ml, 11 mmol) was then slowly added to the flask. The system was allowed to stir until all of the colour had disappeared. Diethyl ether (50 cm³) was added to the flask to ensure that all of the phosphines produced were dissolved. A sample of the resulting solution was collected in a sealed NMR tube and analysed by ³¹P {¹H} NMR.

Table 1

Example	Ratio butyllithium:TMEDA: o-xylene	T °C	time (hrs)	% conversion to disubstituted product (re: o-xylene)	ratio di : mono product
1	2 : 2 : 1	25	96	32	0.7
2	2.5 : 2.5 : 1	25	96	31	0.77
3	2.5 : 2.5 : 1	60	12	32	1.60
4	2.5 : 2.5 : 1	40	6	39	1.08
5	2.5 : 2.5 : 1	60	6	34	0.85
6	3 : 2 : 1	60	6	57	1.50
7	4 : 4 : 1	25	96	57	1.86
8	4 : 2 : 1	25	96	69	2.3

Table 2

	% conversion (mono)	% conversion (di)
solid	5	38
liquid	30	8

Table 3

Example	Ratio BuLi/BuOK/ Xylene	Potassiation reaction time and temperature (°C)	% Conversion di substituted product
10	2.2 : 2.2 : 1	1 hour, room temp. (RT)	40
11	3 : 3 : 1	1 hour at RT + 1 hr @ 60 °C	45
12	2.2 : 2.2 : 1	1 hour @ 80 °C	44
13	3 : 3 : 1	1 hour @ 80 °C	49

Example 14

- 10 O-xylene (3.3 cm³, 270 x 10⁻²mol) was added by syringe to a suspension of potassium t-butoxide (6.6 g, 5.88 x 10⁻²mol) in 40/60 petroleum ether (150 cm³). 42 cm³ of a 1.4 M solution of butyl lithium in hexane was added to the ether mixture, with cooling. The

mixture was stirred overnight to complete the reaction which resulted in an orange-red solid which was the dipotassio-o-xylene. The mixture was filtered and the solid washed with further petroleum ether.

The dipotassio derivative was then suspended in diethyl ether (150 cm³) at -78 °C and
5 di-*t*-butylchlorophosphine (10 g, 5.54 x 10⁻²mol) was added by syringe. The mixture was stirred for an hour, brought slowly to room temperature and then filtered. The filtrate was hydrolysed by addition of 1 cm³ of degassed water and dried. The yield (based on o-xylene) was 94%. Analysis by ³¹P {¹H} NMR showed that the yield of α,α'-bis(dit-butylphosphino)o-xylene was 80%, whilst that of the monophosphine was 10%.

10 Example 15

Potassium-*tert*-butoxide (3.99 g, 35.6 mmol), heptane (150 cm³) and o-xylene (1.47 cm³, 12 mmol) were added to a 500 ml round bottomed flask. To this stirred solution *n*-butyllithium (14.45 cm³, 36.1 mmol, 2.5M in hexane) was added slowly and the flask heated to 80 °C for 1 hour. An orange/red precipitate formed which was separated by
15 filtration and washed with pentane (3 x 100cm³) before being resuspended in diethyl ether (100 cm³) which had been pre-cooled to -20°C. To this solution maintained at -20°C was added di-*t*-butylchlorophosphine (4.58 cm³, 24.1 mmol) and the reaction allowed to warm to room temperature before being stirred overnight. The in-situ reaction yield of α,α'-bis(dit-butylphosphino)o-xylene as determined by ³¹P NMR was 64%.

20 Example 16

The reaction described in Example 15 was repeated except that the orange-red precipitate of dipotassio-o-xylene was resuspended in 100 cm³ of diethyl ether which had been pre-cooled to 0 °C and the solution was maintained at 0°C while di-*t*-butylchlorophosphine (4.58 cm³, 24.1 mmol) was added. The reaction was allowed to warm to room temperature
25 before being stirred overnight. The in-situ reaction yield of α,α'-bis(dit-butylphosphino)o-xylene as determined by ³¹P NMR was 74.9%.

Example 17

The reaction described in Example 15 was repeated except that the orange-red precipitate of dipotassio-o-xylene was resuspended in 100 cm³ of diethyl ether at room temperature
30 and the solution was maintained at room temperature while di-*t*-butylchlorophosphine (4.58

cm³, 24.1 mmol) was added. The reaction was stirred overnight. The in-situ reaction yield of α,α' -bis(dit-butylphosphino)-o-xylene as determined by ³¹P NMR was 75.5%.

Example 18

Sodium-*tert*-butoxide (17.29 g, 180 mmol), *N,N,N',N'*-Tetramethylethylenediamine (27.1
5 cm³, 180 mmol), heptane (100 cm³) and *o*-xylene (7.32 cm³, 60 mmol) were added to a 1
litre round bottomed flask. To this stirred solution *n*-butyllithium (72.0 cm³, 180 mmol, 2.5M
in hexane) was added slowly and the flask heated to 60 °C for 2 hours. An orange
precipitate formed which was separated by filtration and washed with pentane (100cm³)
before being resuspended in pentane (100 cm³). To this was added
10 di-*t*-butylchlorophosphine (22.75 cm³, 120 mmol) and the reaction stirred at room
temperature overnight. The reaction was quenched by the addition of degassed /
deionised water (50 cm³) when two clear layers formed and the pentane layer (upper
layer) was isolated. The solvent was removed in-vacuo and the product purified by
washing with cold methanol (-10C). Yield (20.0g, 84.6%).

Claims

1. A method of manufacturing a compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$, in which each R is independently a pendant, optionally substituted, organic group through which the group is linked to tertiary carbon atom C; L^1 , L^2 are independently a linking group
 - 5 selected from an optionally substituted lower alkylene chain connecting the respective phosphorus atom to the group X and X is a bridging group comprising an optionally substituted aryl moiety to which the phosphorus atoms are linked on available adjacent carbon atoms comprising:
 - i) reacting together a compound of formula $H-L^1-X-L^2-H$ with an organometallic compound
 - 10 to form an intermediate compound of formula $M-L^1-X-L^2-M$, where M is an alkali metal atom,
 - ii) reacting said intermediate compound with a compound of formula $(R_3-C)_2P-A$, where A is a halogen atom, to form said compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$.
2. A method as claimed in claim 1, wherein M is a Group IA metal selected from sodium,
 - 15 potassium or lithium.
3. A method as claimed in claim 1 or 2, wherein R^m-M is generated in situ by an exchange reaction between R^m-Li and potassium or sodium alkyl oxide.
4. A method as claimed in any of claims 1 - 3, wherein the reaction between R^mM and $H-L^1-X-L^2-H$ is carried out in the presence of a basic compound which is capable of forming
 - 20 a complex with the metal.
5. A method as claimed in claim 4, wherein said basic compound comprises tetramethylethylenediamine.
6. A method as claimed in any preceding claim, wherein said compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$ is selected from the group consisting of
 - 25 α,α' bis(di-t-butylphosphino)o-xylene (also known as 1,2 bis (di-t-butylphosphinomethyl) benzene), bis (di-t-neopentyl phosphino) - o - xylene or bis 1, 2 (di-t-butyl phosphino) naphthalene.

7. A catalyst suitable for catalysing the carbonylation of ethylene comprising compound of general formula $(R_3-C)_2P-L^1-X-L^2-P-(C-R_3)_2$ prepared by a method as claimed in claim 1, and palladium or a compound thereof.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/GB 99/00797

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07F9/50 B01J31/24 C07C67/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07F B01J C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 19434 A (ICI PLC) 27 June 1996 cited in the application see example 4	1-7
A	DE 41 34 772 A (BASF AG) 7 May 1992 see the whole document	1-7
A	WO 96 17856 A (GRIFFITH UNIVERSITY) 13 June 1996 see example 2 and scheme II on page 21	1-7
A	GB 877 592 A (ICI LTED) 13 September 1961 see page 2 and claim 1	1-7

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 May 1999

Date of mailing of the international search report

31/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 99/00797

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9619434 A	27-06-1996	AU 701935 B	11-02-1999
		AU 4309596 A	10-07-1996
		BR 9510249 A	04-11-1997
		CA 2207672 A	27-06-1996
		CN 1171098 A	21-01-1998
		CZ 9701932 A	18-02-1998
		EP 0799180 A	08-10-1997
		HU 77016 A,B	02-03-1998
		JP 10511034 T	27-10-1998
DE 4134772 A	07-05-1992	NONE	
WO 9617856 A	13-06-1996	AU 686918 B	12-02-1998
		AU 4168596 A	26-06-1996
		AU 4168696 A	26-06-1996
		WO 9617855 A	13-06-1996
		CA 2207299 A	13-06-1996
		EP 0796266 A	24-09-1997
		JP 10509957 T	29-09-1998
GB 877592 A		NONE	